

## Lawn maintenance calendar\*

AB-MB-NB-NS-ON-PE-QC-SK BC

| STEPS  | FEB | MAR | APR | MAY | JUNE | JULY | AUG | SEPT | ОСТ | NOTES   |
|--|-----|-----|-----|-----|------|------|-----|------|-----|---|
| Seeding Seeding  |     |     |     |     |      |      |     |      |     | The temperature must be between 15 °C and 20 °C for at least two weeks.                                   |
| Raking   |     |     |     |     |      |      |     |      |     | The ground must no longer be soaked.  |
| Aeration   |     |     |     |     |      |      |     |      |     | Use a lawn aerator to make holes 7 to 10 cm in depth.   |
| Dethatching  |     |     |     |     |      |      |     |      |     | Proceed if thatch is more than 1.25 cm thick.   |
| Lawn repairs (M)   |     |     |     |     |      |      |     |      |     | A temperature of 20 °C to 25 °C<br>during the day and 14 °C to 18 °C<br>at night facilitates germination. |
| White grub control   |     |     |     |     |      |      |     |      |     | Water first and then keep lawn moist for 7 days.  |
| Watering   |     |     |     |     |      |      |     |      |     | Water once a week (for a total of 3 cm of water including rainfall), preferably in the morning.           |
| Mowing   |     |     |     |     |      |      |     |      |     | Cut your grass to a height of 8-10 cm. Avoid mowing more than 1/3 of the total height.                    |
| Feeding  |     |     |     |     |      |      |     |      |     | C-I-L Restore Feed & Seed Lawn<br>Fertilizer is perfect in the spring<br>and fall.                        |
| Chinch bug control   |     |     |     |     |      |      |     |      |     | Do not mistake chinch bug damage with symptoms of drought.  |
| Sod webworm control  |     |     |     |     |      |      |     |      |     | Treat as soon as the first signs appear in late summer or as a preventative treatment in spring.          |
| Overseeding  |     |     |     |     |      |      |     |      |     | The temperature must be over 10 °C for optimal germination.   |
| Weed control   |     |     |     |     |      |      |     |      |     | Remove the whole root. Doing that in the fall prevents weeds from reseeding.                              |
| European crane fly control  * The periods suggested above ma |     |     |     |     |      |      |     |      |     | The treatment must target grub larvae.  |

<sup>\*</sup> The periods suggested above may differ depending on the temperature.